

| Intervention | | |
|---|----------------------|-------------------------|
| Equipment/Medication | New Hampshire | |
| 0.25%-0.9% normal saline (NS) | ALS | AEMT/Paramedic/PIFT |
| 2.5%-10% dextrose in water (D_W) | ALS | AEMT/Paramedic/PIFT |
| Abciximab Drip (Reopro) | | PIFT |
| Acetaminophen (Tylenol) PO | BLS(ORALLY) | BLS |
| Acetaminophen (Tylenol) IV | | PIFT |
| Acetylcysteine (Mucomyst) | | PIFT |
| Adjustable compression dressing after sheath pull | | We wouldn't take these. |
| Adenosine | | Paramedic/PIFT |
| Aggrastat | | SCT |
| Albumin | ALS | PIFT |
| Albuterol (MDI) | | BLS |
| Albuterol nebulizer | ALS | AEMT/Paramedic |
| Alupent | | PIFT |
| Amiodarone | ALS | Paramedic/PIFT |
| Amiodarone Drip | ALS | Paramedic/PIFT |
| Angioseal | | Nope |

| | | |
|--|--------------|---------------------|
| Antibiotics | ALS | PIFT |
| Antifungals | ALS | PIFT |
| Antivirals | ALS | PIFT |
| Argatroban | | |
| Aspirin | BLS (ORALLY) | |
| Arterial lines/sheaths | | |
| Atenolol | | PIFT |
| Ativan (Lorazepam) | ALS | Paramedic/PIFT |
| Atracurium | | Paramedic/PIFT |
| Atropine | ALS | Paramedic/PIFT |
| Atrovent (Ipratropium Bromide) | ALS | AEMT/Paramedic/PIFT |
| Automated Internal Cardioverter/Defibrillator (AICD) | BLS | BLS |
| Baby < 28 days old to higher level of care | | |
| Balloon Pump: Intra Aortic - IABP | N/A | |
| Banana bag | ALS | Paramedic/PIFT |
| Barbituates | ALS | ALS |
| Betamethasone | | |
| BIPAP - In Place > 6 Hrs, and no Setting Changes > 6 Hrs | | Paramedic/PIFT |

| | | |
|--|---------------------------------------|---|
| BIPAP - In Place < 6 Hrs, or Setting Changes < 6 Hrs | | Paramedic/PIFT |
| BIVAD | | |
| Blood products | ALS | PIFT |
| Brethine (Terbutaline) | | PIFT |
| Bretylium/Bretylol | | |
| Brevibloc | | |
| Calcium Chloride | ALS | Paramedic/PIFT |
| Calcium Gluconate | ALS | Paramedic/PIFT |
| Catheter (Foley or Condom) | BLS | BLS |
| Carbamazepine (Tegretol) | | |
| Cardene (Nifedipine) | | |
| Cardiac monitoring | ALS | AEMT can take if patient does not have cardiac diagnosis. Paramedics w/ cardiac diagnosis |
| Cardiac assist device | | |
| Cardizem (Diltiazem) | ALS | Paramedic/PIFT |
| Central line | capped- BLS; infusion/monitoring- ALS | capped- BLS; infusion/monitoring- ALS |
| Chest Tube capped w/o suction | BLS | BLS |
| Chest Tube connected to one-way; Hemlich; Pigtail; DRY | ALS | PIFT |
| Chest tube connected to suction; Atrium; Water-seal; WET | ALS | PIFT |

| | | |
|--|-----|---------------------|
| Cimetidine (Tagamet) | ALS | PIFT |
| Cleviprex (clevidipine) | | PIFT |
| Colostomy bag | BLS | BLS |
| Continuous Bladder Irrigation (CBI) | | |
| CPAP via mask | BLS | BLS |
| CPAP via intubation | ALS | PIFT |
| Cryoprecipitate | | PIFT |
| CVP/venous line/sheath - monitor/measuring | N/A | Nope |
| Crofab | | |
| Demerol - facility MUST supply drug | ALS | PIFT |
| Dexamethasone | ALS | Paramedic/PIFT |
| Diphenhydramine (Benadryl) | ALS | Paramedic/PIFT |
| Dextran | | Nope |
| Dextrose - D50/D10/D5N5/D5W - without additives | ALS | AEMT/Paramedic/PIFT |
| Diazepam (Valium) - we do not carry We do | ALS | Paramedic/PIFT |
| Digoxin (Lanoxin) | | PIFT |
| Dilantin (Phenytoin) | ALS | PIFT |
| Dilaudid (Hydromorphone) | ALS | Paramedic/PIFT |

| | | |
|--------------------------------|--|---|
| Diltiazem (Cardizem) | ALS | Paramedic/PIFT |
| Diprivan (Propofol) | | PIFT |
| Dobutamine | ALS | PIFT |
| Dobutrex | | |
| Dopamine Drip | ALS | PIFT |
| ECG monitoring | ALS | AEMT - if the Pt does not have cardiac monitoring Paramedic/PIFT - w/ cardiac diagnosis |
| Epinephrine | BLS/ALS | Basic: Anaphalixis AEMT - push dose for cardiac arrest. Paramedic/PIFT: Anaphalxvis. |
| Esmolol | ALS | PIFT |
| Frozen fresh plasma (FFP) | ALS | PIFT |
| Famotidine (Pepcid) | ALS | PIFT |
| FemSTOP | | |
| Fentanyl (Sublimaze) | ALS | PIFT |
| Fibrinolytics | I would like SCT with big risk of deteration | PIFT |
| Flumazenil (Romazicon) | ALS | PIFT |
| Fosphentoyin | ALS | PIFT |
| Furosemide (Lasix) | They would give this in the ER/Med/surg then send on a 3 hour transfer w/o foley | |
| Glucagon | ALS | Paramedic/PIFT |
| Glycoprotein 11a/11b inhibitor | | |

| | | |
|--|--------------|---------------------|
| Haldol/Haloperidol | ALS | Paramedic/PIFT |
| Heart pump | SCT TEAM | |
| Heparin Injection | | Paramedic/PIFT |
| Heparin drip <18 units/kg/hr OR < 2,000 Units Per Hour | ALS | Paramedic/PIFT |
| Heparin drip >18 units/kg/hr AND > 2,000 Units Per Hour | ALS | Paramedic/PIFT |
| | | |
| | | |
| Hespan | | |
| Hetastarch | | Who knows |
| High Flow Nasal Cannula | Looking into | |
| Hydralazine | | |
| Hydromorphone (Dilaudid) | ALS | Paramedic/PIFT |
| Jackson-Pratt (JP) drain | | |
| Ipratropium Bromide mixed w/ Albuterol (Duoneb) MDI | BLS | BLS |
| Ipratropium Bromide mixed w/ Albuterol (Duoneb) Inhalation | ALS | AEMT/Paramedic/PIFT |
| Ileostomy with suction | | |
| Inderal | | |
| Insulin - not in PPN/TPN | ALS | PIFT |

| | | |
|---|--------------------------------|---------------------|
| Integrillin drip | | |
| Intracranial pressure (ICP) monitoring | N/A | |
| Intrathecal pain pump (subdural implant) | BLS | |
| Intraosseous (IO) lines | ALS | AEMT/Paramedic/PIFT |
| Intubated | SCT TEAM | PIFT |
| IV fluids - central line or pump | ALS | Paramedic/PIFT |
| IV fluids - with electrolytes or nutrients | ALS | Paramedic/PIFT |
| IV pump | ALS | AEMT/Paramedic/PIFT |
| IV steroids | ALS | Paramedic/PIFT |
| Keppra (Levetiracetam) | ALS | PIFT |
| Ketamine | ALS | Paramedic/PIFT |
| Ketorolac (Toradol) | ALS | Paramedic/PIFT |
| Labetalol | ALS (we do not carry) | PIFT |
| Lactated Ringers (LR) | ALS | AMET/Paramedic/PIFT |
| Lanoxin (Digoxin) | ALS | PIFT |
| Levabuterol (Xopenex) MDI | BLS | BLS |
| Levabuterol (Xopenex) | ALS | PIFT |
| Levophed (Norepinephrine) | ALS | Paramedic/PIFT |

| | | |
|--|-----|---------------------|
| Lidocaine for IO | ALS | AEMT/Paramedic/PIFT |
| Lidocaine cardiac push | ALS | Paramedic/PIFT |
| Lidocaine drip | ALS | Paramedic/PIFT |
| Life vest | BLS | BLS |
| Lopressor (Metropolol) | ALS | Paramedic/PIFT |
| Lorazepam (Ativan) | ALS | Paramedic/PIFT |
| Magnesium drip - NOT pregnancy | ALS | Paramedic/PIFT |
| Magnesium - Eclampsia | ALS | Paramedic/PIFT |
| Mannitol | ALS | PIFT |
| MARK 1 | N/A | N/A |
| | | |
| Mediport (port-a-cath/infuse-a-port) | BLS | |
| Medication - Pain Control Pump (PCA) *not central line, patch/paste | BLS | BLS |
| Meperidine - facility must supply | | PIFT |
| Metoclopramide (Reglan) | ALS | Paramedic/PIFT |
| Metaproterenol | ALS | PIFT |
| Methylprednisolone (Solumedrol) | ALS | Paramedic/PIFT |
| Midazolam (Versed) Injection | ALS | Paramedic/PIFT |

| | | |
|---|--------------------------------------|--------------------------------------|
| Midazolam (Versed) drip | ALS | Paramedic/PIFT |
| Milrinone | | |
| Morphine Injection | ALS | Paramedic/PIFT |
| Morphine drip | ALS | Paramedic/PIFT |
| Mynx device (TR BAND) | | |
| Narcan/Naloxone | Nasal- BLS, injection/drip-ALS | S: Nasal AEMT:IVP Paramedic/PIFT: I' |
| Nasogastric tube (NG/NGT) with or without Suction | Without suction-BLS With suction-ALS | Paramedic/PIFT |
| Natreacor | | |
| Neosynephrine | | |
| Nitroglycerin (Nitro; NTG; Tridil) | BLS: with assistance | AEMT/Paramedic/PIFT |
| Nitroglycerin Drip | ALS | Paramedic/PIFT |
| Nitrous Oxide (Laughing gas) | ALS | AEMT/Paramedic/PIFT |
| Octreotide (Sandostatin) | ALS | PIFT |
| Omeprazole (Prilosec) | ALS | PIFT |
| Odanesteron (Zofran) | ALS | AEMT/Paramedic/PIFT |
| Orogastric tube (OG; OGT) | ALS | Paramedic/PIFT |
| Oxygen - any method except ventilator | BLS | |
| Oxymizer Pendant | | |

| | | |
|--|----------------------------|-----------------|
| Packed Red Blood Cells (PRBCs) | ALS | PIFT |
| Pediatrics - < 4 yo or 40lbs | BLS | Depends on call |
| Pediatrics - psych | BLS | BLS |
| Peripheral IV - with fluid or locked | Locked- BLS With fluid-ALS | |
| Peritoneal Dialysis | | Probably not |
| Phenergan (Promethazine) | ALS | PIFT |
| Phenobarbitol | ALS | PIFT |
| Phenylephrine | | |
| Phenytoin (Dilantin) | ALS | Dilantin |
| Phrenetic Nerve Stimulator | BLS | |
| Plasma | ALS | PIFT |
| Plasmanate | | |
| Platelets | ALS | PIFT |
| Potassium < or = 20mEq in 1L bag of fluid | ALS | PIFT |
| Potassium > 20mEq in 1L bag of fluid | ALS | PIFT |
| PPN with insulin | BLS | BLS |
| PPN without insulin | BLS | BLS |
| Precedex (dexmedetomidine) *sedative for intubation | N/A | N/A |

| | | |
|--|-----------------------|---------------------|
| Pregnancy - for OB concern | Depending on the call | |
| Pregnancy - no OB concern | BLS | BLS |
| Pregnancy - not coming from a facility with Labor & Deliver Services (GEC, ECAQ, Kaiser, Chestertown, Cross, Doctors) | Call someone else | |
| Pregnancy - ectopic | Probably be medic | |
| Pregnancy - Magnesium Sulfate | ALS | Paramedic/PIFT |
| Pregnancy - Oxytocin (Pitocin) | ALS | Paramedic/PIFT |
| Pregnancy - psych | BLS | BLS |
| Pressure dressing - after sheath removal | We won't do these | |
| Procainamide | ALS | Paramedic/PIFT |
| Prochlorperazine (Compazine) | ALS | Paramedic/PIFT |
| Pronestyl | ALS | PIFT |
| Protonix | ALS | PIFT |
| Proton Pump Inhibitor (PPI) | | |
| Quinidine | ALS | PIFT |
| Ranitidine (Zantac) | ALS | PIFT |
| Remdesivir (Veklury) | ALS | PIFT |
| Sodium Bicarbonate | ALS | Paramedic/PIFT |
| Sodium Chloride ≤ 0.9 | ALS | AEMT/Paramedic/PIFT |

| | | |
|---|---|----------------------|
| Steroids | ALS | PIFT |
| Succinylcholine | N/A | Perhaps someday |
| Swan Ganz Catheter (PA Catheter) | | |
| Tranexamic Acid (TXA) | ALS | Paramedic/PIFT |
| Telemetry monitoring (TELE) | ALS | |
| Terbutaline Sulfate | ALS | PIFT |
| TBA | | |
| Total Parenteral Nutrition (TPN) | BLS | BLS |
| Tracheostomy (TRACH) | No vent, w/o suctioning- BLS No vent w/suctioning or vented- ALS | |
| Transcutaneous Pacing (TCP) - pads on skin | ALS | Paramedic/PIFT |
| Transvenous Pacemaker (TVP) - wired through skin | ALS | Paramedic/PIFT |
| Urostomy | BLS | BLS |
| Ventricular Assist Device (RVAD/LVAD) | BLS | BLS |
| Vecuronium | N/A | Hopefully someday |
| Ventilator | | PIFT W/ 2nd Provider |
| | | |
| | | |
| | | |

| | |
|--|--|
| | |
| | |

w/o complicated settings

w/o complicated settings

